



## **Geophysical Terranes of the Great Basin and Parts of Surrounding Provinces**

By Jonathan M.G. Glen, Edwin H. McKee, Steve Ludington, David A. Ponce, Thomas G. Hildenbrand, and Melanie J. Hopkins

Open-File Report 2004-1008

2004

U.S. Department of the Interior  
U.S. Geological Survey

---

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

<sup>1</sup> U.S. Geological Survey, 345 Middlefield Road, MS 989 Menlo Park, CA 94025

## TABLE OF CONTENTS

INTRODUCTION .....	1
Physiography.....	1
Potential Field Geophysics .....	2
GRAVITY DATA .....	3
MAGNETIC DATA .....	4
FILTERING AND DERIVATIVE METHODS.....	5
Basement Gravity.....	5
Magnetic Potential (pseudogravity).....	7
Maximum Horizontal Gradients .....	8
POTENTIAL FIELD MAPS .....	8
GEOPHYSICAL TERRANE TABLE .....	9
Generalized Source Rock.....	9
Tectonic Setting .....	11
Provinces.....	12
Scale of Geophysical Features .....	13
DISCUSSION .....	14
ACKNOWLEDGMENTS .....	15
REFERENCES CITED.....	15
<b>Error! Bookmark not defined.</b>	

## FIGURES

Figure	1	Physiographic province map of the Great Basin.
	2	Basement gravity terrane map of the Great Basin.
	3	Magnetic potential (pseudogravity) terrane map of the Great Basin.
	4	Regional isostatic gravity and pseudogravity field maps of the Great Basin.

## TABLES

Table	1	Geophysical features of the Great Basin and parts of surrounding provinces.
	2	Physiographic provinces and their geologic and geophysical character.
	3	Generalized source rock categories .....10
	4	Tectonic setting categories .....11

## INTRODUCTION

This study of geophysical terranes within and surrounding the Great Basin of the western United States (fig. 1A) integrates geophysical and geologic data to provide new insights on basement composition and structure at local, intermediate, and regional scales. Potential field (gravity and magnetic) studies are particularly useful to define the location, depth, and extent of buried basement sources and fundamental structural or compositional boundaries. They especially serve in imaging the subsurface in areas of extensive Cenozoic cover or where surface outcrops may be detached from the deeper crust. Identifying buried compositional or structural boundaries has applications, for example, in tectonic and earthquake hazard studies as they may reflect unmapped or buried faults. In many places, such features act as guides or barriers to fluid or magma flow or form favorable environments for mineralization and are therefore important to mineral, groundwater, and geothermal studies. This work serves in assessing the potential for undiscovered mineral deposits and provides important long-term land-use planning information. The primary component of this report is a set of geophysical maps with anomalies that are labeled and keyed to tables containing information on the anomaly and its source. Maps and data tables are provided in a variety of formats (tab delimited text, Microsoft Excel, PDF, and ArcGIS) for readers to review and download. The PDF formatted product allows the user to easily move between features on the maps and their entries in the tables, and vice-versa. Our goal in highlighting these anomalies is to stimulate thought and research about crustal features of the Great Basin. While we do not offer comprehensive interpretation of every gravity and magnetic feature in the Great Basin, we hope this product will serve as a useful spatial catalog of those features.

### **Physiography**

The Great Basin is a broad hydrologically closed region spanning much of the western United States (fig. 1A). It forms part of the Basin and Range Province (Figure 1B), which is characterized by late Cenozoic regional extension ranging from Mexico nearly to Canada and California to Texas. This extension has been accompanied by normal and detachment faulting, crustal block uplift, downdrop, rotations and tilting, crustal thinning and ductile flow, and

widespread magmatism. Crustal blocks expose a range of variably deformed pre-Cenozoic rocks as well as early and middle Cenozoic volcanic rocks.

The study area (fig. 1A) includes the Great Basin and parts of ten adjacent physiographic provinces (Cascade Mtns, Columbia Plateaus, Snake River Plain, Northern Rocky Mtns, Sierra Nevada, Great Valley, Coast Ranges, Mojave Desert, Colorado Plateaus, and Middle Rocky Mtns). The Great Basin is sharply defined on its northern, western and eastern margins where it is bound by the Snake River Plain, Sierra Nevada, and Colorado Plateaus, respectively. Northwestern, northeastern and southern boundaries, however, are less well defined. Definitions of the Great Basin and its surrounding provinces used in this report largely follow those of Fenneman and Johnson (1946).

### **Potential Field Geophysics**

Several excellent reviews of magnetic methods have been published by Grant (1985a,b), Reford (1980), Hinze and Zietz (1985), Blakely (1995), and Gunn and Dentith (1997). Similarly, Simpson and others (1986), Simpson and Jachens (1989), Jachens and others (1989), and Blakely (1995) provide informative reviews of gravity methods. These geophysical techniques, which allow imaging of subsurface structure, are particularly important in the Great Basin where more than 75% of the surface is covered by Cenozoic deposits. Gravity and magnetic anomalies occur due to lateral contrasts in rock density and magnetic properties (induced and remanent magnetizations), respectively. Rock-property contrasts may occur within a rock unit, such as resulting from gradual lateral facies changes or heterogeneous alteration, or at geologic structures such as faults, folds, or contacts. The geometry and depth to sources, the character of the geomagnetic field, and the rock properties of sources all determine the character of the associated potential field anomalies. Despite the complexity of potential fields and their sources, gravity and magnetic data can be used to resolve the geometry and origin of sources, particularly when combined with other geologic constraints such as the regional tectonic models, surface geology, and seismic data.

## GRAVITY DATA

Gravity data were compiled from a variety of sources reduced and gridded (Hildenbrand and others, 2000) to produce the various gravity and derivative maps shown in this report. Gravity maps, derived from these data, reflect anomalies that may arise from contrasts in density due to contacts between different rock units, partial melting, or phase transitions. Generally, long-wavelength anomalies with smooth gradients originate from sources at depths greater than sources of short-wavelength anomalies that display steep gradients. While short-wavelength anomalies must arise from sources at shallow depths, long-wavelength anomalies, could arise from shallow, thin sources that have gently sloping sides.

In order to produce a gravity map reflecting lateral variations in density in the crust, raw gravity measurements were reduced using standard gravity reduction methods (Dobrin and Savit, 1988; Blakely, 1995). These reductions remove the effects of elevation, topography, total mass, rotation, and ellipsoidal shape of the Earth and yield the complete Bouguer gravity anomaly (CBA). Although the CBA reveals lateral density variations at short wavelength scales, it does an inferior job isolating longer wavelength features since these are often masked by broad anomalies due to deep crustal roots that isostatically compensate topographic loads. The isostatic correction attempts to remove effects of compensating masses.

Despite its name, an isostatic anomaly does not necessarily reflect the state of isostatic equilibrium. In this study, we are most interested in those cases where anomalies arise from density inhomogeneities in the crust. Nonetheless, it should be noted that an isostatic anomaly may reflect areas out of isostatic balance, either dynamically or by means of elastic support. For example, because the isostatic correction ignores effects of lithospheric strength it may not accurately account for regional fields associated with topographic loads. Thus, in areas where the lithosphere is exceptionally strong, topographic loads can be supported regionally, and compensation distributed over the area of elastic flexure rather than being locally compensated.

A general correlation between topography and the regional Bouguer gravity field indicates that across the Great Basin the crust is in isostatic equilibrium. A common misconception is that the depth of compensation corresponds to the Moho. Because Bouguer gravity does not correlate to crustal thickness (Eaton and others, 1978) and gravity effects due to variations in the crustal

thickness versus lithospheric thickness differ by an order of magnitude (i.e., asthenosphere/lithosphere density contrast is  $\sim 0.08$  g/cc and between crust and lithosphere it is  $\sim 0.4$  g/cc), compensation likely occurs at mid-crustal depths. In the Great Basin, where heat flow is high and extension is large, isostatic compensation is probably achieved over relatively short lateral distances (e.g. 150 km), and within relatively shallow crustal depths perhaps within the upper 20 km (Eaton and others, 1978).

Although the close correspondence of topography and Bouguer gravity suggests compensation is achieved largely by an Airy-Heiskanen mechanism, some regions, such as the Rocky Mountains (Eaton and others, 1978, Woollard, 1972), are best fit with a Pratt model that achieves compensation through lateral density variations. This is indicated by crustal density variations inferred from seismic compressional wave velocity data. It has been suggested that isostatic equilibrium and extension in the Great Basin are likely accommodated by shallow crustal intrusion or ductile flow, especially in areas such as core complexes where the crust has undergone significant stretching and thinning (Thompson and McCarthy, 1990).

## **MAGNETIC DATA**

Magnetic data were derived from a compilation of statewide compilations (Hildenbrand and others, 2000). Surveys have all been continued to a common reference level of 0.305 km above ground surface, gridded, and filtered to produce the magnetic maps shown in this report. Variations in the magnetic field arise from contrasts in the magnetic properties of rocks. These contrasts can be due to a number of different sources including crustal structures, metamorphism and alteration, variations in remanent magnetization, and variations in the concentration and type of magnetic minerals.

Generally, the most significant contributions to crustal anomalies arise from magnetite, one of the most common, strongly magnetic minerals in the crust. Furthermore, because the magnetic susceptibility of magnetite dramatically drops at temperatures above the Curie point ( $580^{\circ}$  for pure magnetite), the portion of crust most responsible for variations in the magnetic field is that

which lies above the Curie isotherm. This probably coincides roughly with the Moho, though in areas of high heat flow, like the Great Basin, it may occur at significantly shallower depths.

Although the magnetic field strength depends on both induced and remanent crustal magnetization, it is often assumed that the remanent component is negligible. This is because remanence is often low enough to ignore or because remanent components are often aligned close to the induced field component. In general, this is supported by the character of many magnetic anomalies. For example, in the northern hemisphere, anomalies often have relatively weak minima that lie to the north of their maxima counterparts (note that magnetic sources generally display bipolar anomalies). An important effect on the character of geophysical anomalies is the depth to the source. The shallower the depth to a body, the higher the amplitude, the shorter the wavelength, and the sharper the gradients of its anomaly. Generally, magnetic highs arise from mafic igneous and crystalline basement rocks, whereas lows arise from felsic igneous, sedimentary, or altered basement rocks. Igneous outcrops not associated with high-amplitude magnetic anomalies might be thin or contain low concentrations of primary magnetic minerals, or have lost them due to alteration.

Aeromagnetic anomalies in most of the Great Basin have been found to arise from Precambrian metamorphic, Mesozoic granitic and gabbroic, Tertiary calc-alkaline volcanic and intrusive, and Tertiary basaltic rocks (Blakely, 1988). Much of the Precambrian basement in the Great Basin is weakly magnetic, in contrast to the magnetic basement of the Colorado Plateaus (Mabey and others, 1978).

## **FILTERING AND DERIVATIVE METHODS**

### **Basement Gravity**

An iterative gravity inversion method (Jachens and Moring, 1990), used to determine the depth to pre-Cenozoic basement and the thickness of Cenozoic basin deposits, was applied to the Great Basin to obtain a basement gravity map (Figure 2), a by-product of the depth-to-basement process. Basement gravity, which is the isostatic gravity with the effects of Cenozoic basins

removed, reflects lateral density variations in pre-Cenozoic basement rocks and is particularly useful for defining pre-Cenozoic structures and crustal geophysical terranes.

Isostatic gravity anomalies were used during this inversion process because they enhance or reflect shallow- to mid-crustal sources within the Earth by removing long-wavelength variations in the gravity field inversely related to regional topography (Simpson and others, 1986). The Basement gravity inversion process depends on their being a significant contrast in density between the usually dense basement rocks and any overlying Cenozoic deposits. While this is true for much of the Great Basin, Cenozoic mafic volcanic rocks can have densities similar to those of their underlying basement rocks. This is a problem, particularly in the northwestern-most part of the Great Basin that is blanketed by middle Miocene mafic lava flows. As a result, the inversion process was not applied in areas where the thickness of overlying volcanic rocks could not be determined. The boundary within which the depth-to-basement calculation was applied is shown in Figure 2. Outside this boundary, we show isostatic gravity values.

The depth-to-basement method separates the gravity field into two components: the gravity field caused by pre-Cenozoic basement and the gravity field caused by overlying younger basin deposits. An initial basement gravity field is determined by using only stations located on pre-Cenozoic basement outcrops. The initial basement gravity field is approximate because stations located on basement are influenced by the gravity effect of low-density deposits in nearby basins, especially for those stations near the edge of the basins. The difference between the isostatic gravity and basement gravity fields provides the first estimate of the basin gravity field, which is inverted to provide the first estimate of the basin shape. The gravitational effect of the basins is subtracted from each station located on basement and a new and improved basement gravity field is determined. This process is repeated until successive iterations converge. Inversion of the final basin gravity field yields the final estimate of the depth to pre-Cenozoic basement. The density of basement rocks is allowed to vary horizontally, whereas the density of basin-filling deposits increases with depth according to a density-depth relationship defined by Jachens and Moring (1990).

A number of limitations are inherent in this method, including uncertainties that relate to: the gravity data coverage, especially for stations on basement outcrops; the density-depth function; accuracy and scale of the geologic mapping; simplifying assumptions regarding concealed



geology; and the distribution of basement outcrops. A more detailed discussion of the limitations and accuracy of the method were provided by Jachens and Moring (1990).

### **Magnetic Potential (pseudogravity)**

Crustal magnetism differs from and is more complex than gravity, which varies due simply to the crustal density distribution. Magnetism varies because of differences in both the concentration and type of magnetic minerals within the crust (analogous to the relation between density and gravity), and crustal remanent magnetization. Furthermore, because crustal magnetization is seldom vertical, except at the magnetic poles, anomalies are asymmetric and not centered over their sources. In addition, unlike gravity, crustal remanent magnetism has a depth limit set by the Curie temperature isotherm, the temperature above which remanent magnetization does not exist. Magnetic data also tend to highlight shallower features than gravity, because magnetic field strength attenuates more significantly with distance to the source than does gravity.

Because of this complexity of magnetic anomalies they are typically more difficult to interpret. The pseudogravity or magnetic potential transformation (Baranov, 1957; Blakely, 1995) removes asymmetry of anomalies, by centering them over their sources, and allows for a more accurate estimate of the extent of source bodies. In addition, it helps highlight regional magnetic features masked by high-frequency anomalies.

Because the magnetic and gravity potentials are related by a directional derivative, thus the total magnetic field can be transformed into an equivalent gravity field. Magnetic potential, or pseudogravity, maps are produced by the transformation of the magnetic field into the equivalent gravity field assuming a density distribution equal to the magnetization distribution (Baranov, 1957). The ratio between magnetization and density is held constant and remanent magnetization is assumed to be either negligible or in the same direction as the Earth's magnetic field. This process amplifies long wavelengths (deeper sources) at the expense of short wavelengths (shallow sources). In addition, because gravity anomalies have their steepest gradients approximately over the edges of their causative sources, especially for shallow sources, the magnetic potential map can be used to approximate the edges of magnetic sources (Blakely, 1995).

## **Maximum Horizontal Gradients**

To better define the edges of geophysical sources and to help derive geophysical lineaments and terranes, the amplitudes of the maximum horizontal gradients (AMHG) of both gravity and magnetic data were computer generated. A technique described by Blakely and Simpson (1986) was used to calculate the AMHG. Because the AMHG tend to lie over the edges of bodies with near vertical boundaries (Cordell and McCafferty, 1989; Grauch and Cordell, 1987), they are useful at estimating the extent of buried sources. AMHG were derived for both previously described basement gravity and magnetic potential maps. Because these maxima reflect abrupt lateral changes in the density or magnetization of the underlying rocks, they were used to aid in defining the boundaries of geophysical terranes shown in Figures 2 and 3.

## **POTENTIAL FIELD MAPS**

Gravity and magnetic lineations (shown in figs. 2 and 3, respectively) were derived with the aid of the AMHG method described above. Geophysical terranes are based in part on the AMHG-derived boundaries and on geophysical fabric. Areas, for example, that display a consistent trend or wavelength of anomalies, in contrast to their surroundings, were defined as distinct geophysical terranes that may represented discrete crustal blocks having similar physical properties or sources. Gravity and magnetic terrane maps (figs. 2 and 3) were created by visual inspection of gravity, magnetic, and derivative geophysical maps, and by drawing polygons around similar geophysical areas using derived lineaments as a guide to locating terrane boundaries. In addition to geophysical terranes, we have also included a number of lineations. A few of these are defined as geophysical features listed in table 1, while others occur on figures 2 and 3 simply as unlabeled features that are intended to highlight the geophysical fabric.

## **GEOPHYSICAL TERRANE TABLE**

The geophysical terrane table (table 1) lists geophysical terranes that occur on the gravity and magnetic terrane maps (figs. 2 and 3). The table is organized to allow the user to identify features in the table that occur on the maps, and to move between maps and table. Terrane names contain (from left to right): a two letter (uppercase) code identifying the state in which they mostly occur, a single letter (lowercase) code identifying whether the feature is a gravity (g) or magnetic (m) terrane, and a two digit number indicating the feature number. Geologic and geophysical references provided in the table point the user to an example of work pertaining to some part of the anomaly or to its presumed source rock, and should not be considered a complete list of pertinent or historical citations. We refer the reader to references contained within the cited publications for further background. Several of the table columns are specifically defined such that they may be used as search terms or as tools for sorting the table based on the terrane characteristics (These include: Generalized Source Rock, Province, Tectonic Setting, and Scale). A brief discussion of these terms is provided below. These search fields are inherently simplified and may not adequately explain the character, especially of diverse terranes.

### **Generalized Source Rock**

Five primary rock categories (sedimentary, volcanic, intrusive, metamorphic, and basement) are used in conjunction with five secondary rock categories (silicic, mafic, ultramafic, carbonate, and siliceous) to provide thirteen categories (table 3) to describe the generalized source rock. Note that in some cases, the choice between terms is arbitrary. For example, basement and metamorphic rock categories overlap, and in many cases are interchangeable. Here, 'Metamorphic' is generally used in the Coast Ranges and Klamath Mountains, and 'Basement' in the Great Basin and Colorado Plateaus. In areas where several different source rocks may be present, multiple rock type terms have been used. 'Basement' is generally used throughout this report to refer loosely to dense, crystalline, and usually Precambrian rocks. This is in contrast to its use in the term 'basement gravity' (e.g., fig. 2), which considers basement as pre-Cenozoic rocks assumed to be dense, crystalline rocks of many types. Note that in places where source

rocks are entirely covered, the inferred source rock type is generally inferred from the gravity and magnetic character of the terrane, and may not be unique. For example, a gravity and magnetic low that could be inferred as due to silicic basement might instead be due to depressed basement that has no lateral variation in basement composition.

Table 3. Generalized source rock categories

Primary	Secondary	Usage
sedimentary	carbonate	Generally used to refer to dense sedimentary rocks (e.g. limestone, dolomite).
	siliceous	Generally used to refer to non-carbonate sedimentary rocks (e.g. sandstone, siltstone, argillites, cherts).
volcanic	silicic	Generally used for non-magnetic volcanic rocks. An exception to this is in areas of silicic tuff. Generally includes intermediate composition rocks (e.g. andesites).
	mafic	Generally used for magnetic volcanic rocks.
intrusive	silicic	Generally used for low density or non-magnetic intrusive rocks. Generally includes intermediate composition rocks (e.g. dacites).
	mafic	Generally used for dense or magnetic intrusive rocks. Note that some granitic rocks can be dense.
	ultramafic	Generally used for dense or magnetic intrusive rocks.
metamorphic	silicic	Generally used for low density or non-magnetic (e.g. quartzose) metamorphic rocks.
	mafic	Generally used for dense and magnetic metamorphic rocks.
	ultramafic	Generally used for dense and magnetic metamorphic rocks.
basement	silicic	Generally used for relatively low density or non-magnetic basement rocks.

	mafic	Generally used for very dense and magnetic basement rocks.
	carbonate	Generally used for dense and non-magnetic basement rocks.

## Tectonic Setting

The ‘Tectonic Setting’ category describes the tectonic setting associated with the development of the geophysical feature. In cases where a terrane formed from several geologic events or during multiple stages, or straddled different tectonic settings, multiple categories of ‘Tectonic Setting’ were used. As a result, both ancient and recent settings may be listed. Nonetheless, the most representative setting controlling the character of the feature is given (e.g., the ‘Tectonic Setting’ of Salinian granites in the Coast Ranges is given as both ‘Batholith’ and ‘Transcurrent’ because the shape of the block subsequent to the rocks having formed as a batholith was modified by transcurrent strike-slip motion). In cases where multiple terms are given, the setting considered primarily responsible for the feature is listed first. When there is doubt as to the origin of the source, the present tectonic setting is given (e.g. ‘Extension’ applies to sources within the Great Basin with no known tectonic origin). Table 4 lists ‘Tectonic Setting’ categories that include eleven terms (extension, transcurrent, compression, stable crust, subduction, accretion, uplift, depression, hotspot, batholith, continental margin). The use of ‘Subduction’ to describe the tectonic setting of terranes within the Great Basin applies to Mesozoic intrusive rocks mainly in the western Great Basin (Walker Lane Belt) and Oligocene to middle Miocene volcanism that occurred throughout the Great Basin due to shallowing and subsequent steepening of the subducting Farallon slab.

Table 4: Tectonic setting categories

Tectonic setting	Usage
extension	Generally applies to entire Great Basin with Basin and Range type extension and to even greater extended terranes. Also used to describe back-arc spreading as seen in the Oregon highlands.

transcurrent	Refers to areas subjected to significant strike-slip deformation. Generally applied to right-lateral displacements in Coast Ranges and the Walker Lane.
compression	Used in the region east of the Idaho Batholith and north of the Snake River Plain -- an area in the Rocky Mountains Foreland Thrust Belt.
stable crust	Largely applies to the Colorado Plateaus region and refers to areas of weakly deformed flaying Mesozoic and Paleozoic rocks.
subduction	Applies to major batholiths and to extensive magmatism in the Great Basin thought to relate to shallowing of the subducting Farallon Plate in the later part of the Cenozoic.
accretion	Used extensively in California in the Coast Ranges, Klamath Mountains, and extending to the western Sierra Nevada. These areas are often also associated with transcurrent tectonic setting.
uplift	Used for blocks of basement uplifted on high-angle faults or on flexures. Areas include core complexes and the Colorado Plateaus.
depression	Used largely for isolated basins in the California Coast Ranges. Also associated with areas of pull-apart tectonics. Can also apply to depressed basement.
hotspot	Used to refer to magmatism and fracturing associated with the ancestral Yellowstone hotspot.
batholith	Used to refer to large granitoid intrusions associated with the Sierra Nevada, Salinian Block, and Idaho Batholiths and to fragments of these such as may occur eastern California and west-central Nevada.
continental margin	Refers to Paleozoic and Mesozoic continental margin in the Great Basin. Defined essentially by the edge of the continental shelf.

## Provinces

Geophysical terranes within twelve physiographic provinces, that include the Great Basin and parts of its surrounding provinces (Cascade Mtns., Coast Ranges, Colorado Plateaus, Columbia Plateaus, Great Valley, Klamath Mtns., Middle Rocky Mtns., Mojave Desert, Northern Rocky Mtns., Sierra Nevada, Snake River Plain) are described in this report (figs. 2 and 3, table 1). Figure 1 shows the extent of these provinces (note that the province boundaries used here largely follow those of Fenneman and Johnson (1946), and table 2 gives their physiographic, geologic and geophysical descriptions.

In places, province boundaries are poorly defined, for example, at the boundary between the Great Basin and Columbia Plateaus Provinces. This boundary, which is not well expressed physiographically, represents one of the few differences between the province map used here and that of others (e.g. Fenneman and Johnson, 1946). Terms used under the 'Generalized Tectonic Setting' column refer to the same terms used in Table 4 and described above.

## **Scale of Geophysical Features**

At the broadest scales, geophysical terranes can reflect major deep-seated crustal discontinuities such as transform, accommodation, or shear zones, ancient continental margins, failed rifts, accretionary belts, or magmatic arcs. At local scales, they can reflect, for example, individual faults or intrusive bodies. Below is a description and definition of terrane scales that appear in table 1.

### **Regional-scale Geophysical Provinces**

Regional scale terranes are considered to constitute very large regions, extending from hundreds to thousands of kilometers, that may consist of an assemblage of smaller-scale features that share a common character in contrast to surrounding regions (e.g. the terrane may define a zone of consistent geophysical fabric). A regional-scale feature may reflect a region of common tectonic or magmatic history and it may be bound by deep crustal to subcrustal faults. Some examples include volcanic plateaus, broad shear zones, and broad and coherent crustal blocks bound by deep crustal faults. Identification of regional-scale features is aided by standard and long-wavelength geophysical maps, fabric analysis, contrasts of dominant frequencies, and contrasts in mean gravity and magnetic values.

A description of the regional geophysical expression of the Great Basin and each of its surrounding provinces is given in table 2. The broadest expression in regional gravity and magnetic maps of the study area are reflected in figure 4, which show outlines that roughly mimic the shape and extent of the entire Great Basin. These largest of terranes appear in table 1 as the first two entries.

The Great Basin boundary has relatively sharply defined geophysical boundaries, though these, in places, crosscut the physiographically and geologically defined boundaries of the region. To the north it contrasts with the high basement gravity and magnetic character of the Snake River Plain, although the gravity high over the northern Great Basin merges with that of the Snake River Plain. To the west, the high frequency magnetic and moderate gravity highs of the western

Great Basin about the prominent northwest-trending magnetic high and gravity low of the eastern Sierra Nevada Province. At its east edge, the Great Basin is rimmed by the Colorado Plateaus. Though its contact is not particularly clear in the basement gravity, the generally low magnetic fields over the Great Basin contrast sharply with the high magnetic terrain of the Colorado Plateaus. More ambiguous occur to the northeast, northwest, and south where the Great Basin meets with Middle Rocky Mountains, Columbia Plateaus, and Mojave Desert Provinces, respectively.

### **Intermediate-Scale Geophysical Features**

Intermediate-scale terranes constitute large coherent anomalies (e.g., crustal rifts or sutures, structural basins or ranges, or batholiths) that extend on the order of tens to hundreds of kilometers. Identification of these regional-scale features is aided by standard geophysical maps, long- and short-wavelength maps, and AMHG maps.

### **Local-Scale Geophysical Features**

Local-scale features (reflecting, for example, individual plutons, faults, or calderas) reflect anomalies arising from discrete source bodies that reside in the shallow to mid-crust, and extend over several tens of kilometers. Identification of local-scale features is aided by standard geophysical, long- and short-wavelength, and AMHG maps. Although a few local-scale anomalies are described in this report, a detailed assessment of local-scale anomalies is beyond the scope of this study.

## **DISCUSSION**

This work provides an introduction to geophysical terranes of the Great Basin and surrounding regions, with interpretations of their underlying causes. The aim of this work is to relate geophysical terranes to their geologic domain counterparts, to resolve the nature of transitions



between terranes, and to understand their origin, and their relation to basement structures and composition.

The study spans local-scale (e.g., individual plutons, faults, or calderas), anomalies arising from discrete source bodies that reside in the shallow to mid-crust, and extend over an order of several tens of kilometers to regional-scale features (very large regions, extending from hundreds to thousands of kilometers). Assessment of geophysical terranes was aided by a variety of filtering and derivative methods, and took into account frequency, amplitude, fabric, and gradients of terranes anomalies.

Geophysical maps and data tables of this report are provided in a variety of formats (tab delimited text, Microsoft Excel, Microsoft Word, PDF, and ArcGIS) for readers to review and download. The PDF formatted product contains useful links that allow the user to easily move between features on the maps and their entries in the tables.

## **ACKNOWLEDGMENTS**

We thank George Billingsley, Richard Blakely, Mark Gettings, Bob Jachens, David John, Victor Mossotti, David Piper, Ted Theodore, and Allan Wallace for helpful reviews and discussions, and Barry Moring for assisting in technical preparation of this manuscript.

## **REFERENCES CITED**

Baranov, V., 1957, A new method for interpretation of aeromagnetic maps: pseudo-gravimetric anomalies: *Geophysics*, v. 22, p. 359-383.

Best, M.G., Christiansen, E.H., and Blank, R.H., Jr., 1989a, Oligocene caldera complex and calc-alkaline tuffs and lavas of the Indian Peak volcanic field, Nevada and Utah: *Geological Society of America Bulletin*, v. 101, p. 1076-1090.

Best, M.G., Christiansen, E.H., Deino, A.L., Gromme, C.S., McKee, E.H., and Noble, D.C., 1989b, Excursion 3A - Eocene through Miocene volcanism in the Great Basin of the western United States: New Mexico Bureau of Mines and Mineral Resources Memoir 47, p. 91-133.

Best, M.G., McKee, E.H., and Damon, P.E., 1980, Space-time composition patterns of Late Cenozoic mafic volcanism, southwestern Utah and adjoining areas: American Journal of Science, v. 280, p. 1035-1050.

Blackwell, D.D., 1978, Heat flow and energy loss in the Western United States, *in* Smith, R.B., and Eaton, G.P., eds., Cenozoic tectonics and regional geophysics of the Western Cordillera: Geological Society of America Memoir 152, p.175-208.

Blakely, R.J., 1988, Curie temperature isotherm analysis and tectonic implications of aeromagnetic data from Nevada: Journal of Geophysical Research, v. 93, p. 11,817-11,832.

Blakely, R.J., 1995, Potential theory in gravity and magnetic applications: New York, Cambridge University Press, 441 p.

Blakely, R.H., and Jachens, R.C., 1991, Regional study of mineral resources in Nevada: Insights from three-dimensional analysis of gravity and magnetic anomalies: Geological Society of America Bulletin, v. 103, p. 795-803.

Blakely, R.H., Jachens, R.C., Simpson, R.W., and Couch, R.W., 1985, Tectonic setting of the southern Cascade Range as interpreted from its magnetic and gravity fields, Geological Society of America Bulletin, v. 96, p. 43-48.

Blakely, R.J., Schruben, P.G., and Moring, B.C., 1996, Shallow magnetic lithologies as interpreted from low-altitude aeromagnetic data, in Singer, D.A., An analysis of Nevada's metal-bearing mineral resources: Nevada Bureau of Mines and Geology Open-File Report 96-2, p. 3.1-3.9.

Blakely, R.J., and Simpson, R.W., 1986, Approximating edges of source bodies from gravity or magnetic data: Geophysics, v. 51, p. 1494-1498.

Blank, H.R., Jr., 1988, Basement structure in the Las Vegas region from potential-field data: Geological Society of America, Abstracts with Programs, v. 20, no. 3, p. 144.

- Blank, H.R., Butler, W.C., and Saltus, R.W., 1998, Neogene uplift and radial collapse of the Colorado Plateau - regional implications of gravity and aeromagnetic data: U.S. Geological Survey Bulletin 2158, p. 9-32.
- Bott, M.H.P., and Smithson, S.B., 1967, Gravity investigations of subsurface shape and mass distributions of granite batholiths: Geological Society of America Bulletin, v. 78, p. 859-878.
- Burnett, J.L., and Jennings, C.W., 1982, Bouguer gravity map of California, Chico Sheet: California Division of Mines and Geology Map, scale 1:250,000.
- Byerly, P.E., 1966, Interpretations of gravity data from the central Coast Ranges and San Joaquin valley: California Geological Society of America Bulletin, v. 77, p. 83-94.
- Byers, F.M., Jr., Carr, W.J., Orkild, P.P., Quinlivan, W.D., and Sargent, K.A., 1976, Volcanic suites and related calderas of Timber Mountain-Oasis Valley caldera complex, southern Nevada: U.S. Geological Survey Professional Paper 919, 70 p.
- Cady, J.W., 1975, Magnetic and gravity anomalies in the Great Valley and western Sierra Nevada metamorphic belt, California: Geological Society of America Special Paper 168, 56p.
- Case, J.E., and Joesting, H.R., 1972, Regional geophysical investigations in the central Colorado Plateau: U. S. Geological Survey Professional Paper 736, 31p.
- Chapman, R.H., and Griscom, A., 1980, Coast Ranges, *in* Oliver, H.W., ed., Interpretation of the gravity map of California and its continental margin: California Division of Mines and Geology Bulletin 205, p. 24-27.
- Chapman, R.H., Healey, D.L., and Toxel, B.W., 1973, Bouguer gravity map of California, Death Valley sheet: California Division of Mines and Geology Map, scale 1:250,000.
- Chen, J.H., and Moore, H. G., 1982, Uranium-lead isotopic ages from the Sierra Nevada Batholith, California: Journal of Geophysical Research, v. 87, no. 129-133.
- Cordell, L., and McCafferty, A.E., 1989, A terracing operator for physical property mapping with potential field data: Geophysics, v. 54, p. 621-634.

Craig, S.D., ed., 1992, Walker Lane Symposium: structure, Tectonics and mineralization of the Walker Lane: Geological Society of Nevada, Proceeding., v. 14, April 24, 65p.

Dallmeyer, R.D., Snook, A.W., and McKee, E.H., 1986, Mesozoic-Cenozoic tectonothermal evolution of the Ruby Mountains, east Humboldt Range, Nevada: a Cordilleran metamorphic core complex: Tectonics, v. 5, p. 931-954.

Davis, G.A., and Burchfiel, B.C., 1973, Garlock Fault: An Intracontinental Transform Structure, Southern California: Geological Society of America Bulletin, v. 84, p. 1407-1422.

Dilek, Y., and Moores, E.M., 1995, Geology of the Humboldt igneous complex, Nevada, and tectonic implications for the Jurassic magmatism in the Cordilleran Orogen: Geological Society of America Special Paper 299, p. 229-248.

Dobrin, M.B., and Savit, C.H., 1988, Introduction to Geophysical Prospecting (4th ed.): New York, McGraw-Hill, 867p.

Eaton, G.P., Wahl, R.R., Prostka, H.J., Mabey, D.R., and Kleinkopf, M.D., 1978, Regional gravity and tectonic patterns: Their relation to late Cenozoic epeirogeny and lateral spreading in the western Cordillera, *in* Smith, R.B., and Eaton, G.P., eds., Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152, p. 51-91.

Eckel, E.B., ed., 1968, Nevada Test Site: Geological Society of America Memoir 110, 290 p.

Fenneman, N.M., 1931, Physiography of western United States: New York and London, McGraw Hill Co, 534p.

Fenneman, N.M., and Johnson, D.W., 1946, Map of physical divisions of the United States, Prepared in cooperation with the Physiographic Committee of the Geological Survey, scale 1:7,000,000.

Glen, J.M.G., and Ponce, D.A., 2002, Large-scale fractures related to inception of the Yellowstone hotspot: Geology, v. 30, p. 647-650.

Glen, J.M.G., Hildenbrand, T.G., and Ludington, S., 2000, Insights on the basement of the Great Basin from an integrated geophysical, geologic and geochemical study: Geological Society of America, Abstracts with programs, v. 32, p. A503.

Grant, F.S., 1985a, Aeromagnetics, geology and ore environments; II, Magnetite and ore environments: *Geoexploration*, v. 23, p. 335-362.

Grant, F.S., 1985b, Aeromagnetics, geology and ore environments; I, Magnetite in igneous, sedimentary and metamorphic rocks; an overview: *Geoexploration*, v. 23, p. 303-333.

Grauch, V.J.S., 1996, Magnetically interpreted, granitoid plutonic bodies in Nevada, *in* Singer, D.A., ed., An analysis of Nevada's metal-bearing mineral resources: Nevada Bureau of Mines and Geology Open-File Report 96-2, p. 7.1-7.16.

Grauch, V.J.S., and Cordell, L., 1987, Limitations of determining density or magnetic boundaries from the horizontal gradient of gravity or pseudogravity data: *Geophysics*, v. 52, p. 118-121.

Grauch, V.J.S., Jachens, R.C., and Blakely, R.J., 1995, Evidence for a basement feature related to the Cortez disseminated gold trend and implications for regional exploration in Nevada: *Economic Geology*, v. 90, p. 203-207.

Grauch, V.J.S., Sawyer, D.A., Fridrich, C.J., and Hudson, M.R., 1999, Geophysical framework of the southwestern Nevada volcanic field and hydrogeologic implications: U.S. Geological Survey Professional Paper 1608, 39p.

Griscom, A., 1966, Magnetic data and regional structure in northern California, *in* *Geology of northern California*: California Division of Mines and Geology Bulletin 190, p. 407-417.

Griscom, A., 1973, Bouguer gravity map of California, Redding Sheet: California Division of Mines and Geology Map, scale 1:250,000.

Griscom, A., 1980a, Cascade Range and Modoc Plateau, *in* Oliver, H.W., ed., Interpretation of the gravity map of California and its continental margin, California Division of Mines and Geology, Bulletin 205, p. 36-38.

Griscom, A., 1980b, Klamath Mountains Province, *in* Oliver, H.W., ed., Interpretation of the gravity map of California and its continental margin: California Division of Mines and Geology, Bulletin 205, p. 34-36.

Griscom, A., and Jachens, R.C., 1990, Tectonic implications of gravity and magnetic models along east-west seismic profiles across the Great Valley near Coalinga, *in* Rymer, M.J., and Ellsworth, W.L., eds., The Coalinga, California earthquake of May 2, 1983: U.S. Geological Survey Professional Paper 1487, p. 69-78.

Griscom, A., and Oliver, H.W., 1980, Isostatic gravity highs along the west side of the Sierra Nevada and the Peninsular Ranges batholiths, California: *Eos*, Transactions of the American Geophysical Union, v. 61, p. 1126.

Gunn, P.J., and Dentith, M.C., 1997, Magnetic response associated with mineral deposits, *in* Gunn, P.J., ed., Airborne magnetic and radiometric surveys: AGSO Journal of Australian Geology and Geophysics, v. 17, p. 145-158.

Hanna, W.F., Burch, S.H., and Dibblee, T.W., Jr., 1972, Gravity, magnetics, and geology of the San Andreas Fault area near Cholame, California: U. S. Geological Survey Professional Paper 646-C, p. C1-C29.

Healey, D.L., 1970, Bouguer gravity map of California, Kingman Sheet: California Division of Mines and Geology Map, scale 1:250,000.

Henry, C.D., and Ressel, M.W., 2000, Interpretation of Eocene magmatism, extension, and Carlin-type gold deposits in northeastern Nevada, *in* Lageson, D.R., Peters, S.G., and Lahren, M.M., eds., Great Basin and Sierra Nevada: Geological Society of America Field Guide 2, p. 165–187.

Hildenbrand, T.G., Berger, B., Jachens, R.C., and Ludington, S., 2000, Regional crustal structures and their relationship to the distribution of ore deposits in the western United States, based on magnetic and gravity data: *Economic Geology*, v. 95, p. 1583-1603.

Hinze, W.J., and Zietz, I., 1985, The composite magnetic-anomaly map of the conterminous United States, *in* Hinze, W.J., ed., The utility of regional gravity and magnetic anomaly maps: 52nd annual international meeting of the Society of Exploration Monograph, p. 1-24.

Hutchinson, R.W., and Albers, H.P., 1992, Metallogenic evolution of the Cordilleran region of the Western United States, *in* Burchfiel, B. C., Lipman, P.W., and Zoback, M.L., eds., The geology of North America, The Cordilleran orogen: Conterminous U.S.: Boulder, Colorado, Geological Society of America, v. G-3, p. 629-652.

Jachens, R.C., and Griscom, A., 1985, An isostatic gravity map of California: a residual map for interpretation of anomalies from intracrustal sources, *in* Hinze, W.J., ed., The utility of regional gravity and magnetic anomaly maps: Tulsa, Oklahoma, Society of Exploration Geophysicists, p. 347-360.

Jachens, R.C., and Griscom, A., *in press*, Geologic and Geophysical setting of the 1989 Loma Prieta earthquake, California, inferred from magnetic and gravity anomalies: U.S. Geological Survey Professional Paper.

Jachens, R.C., and Griscom, A., and Roberts, C.W., 1995, Regional extent of Great Valley basement west of the Great Valley, California: implications for extensive tectonic wedging in the California Coast Ranges: *Journal of Geophysical Research*, v. 100, p. 12769-12790.

Jachens, R.C., and Moring, B.C., 1990, Maps of the thickness of Cenozoic deposits and the isostatic residual gravity over basement for Nevada: U.S. Geological Survey Open-File Report 90-404, 15p.

Jachens, R.C., Simpson, R.W., Blakely, R.J., and Saltus, R.W., 1989, Isostatic residual gravity and crustal geology of the United States, *in* Pakiser, L.C., and Mooney, W.D., eds., Geophysical framework of the continental United States: Geological Society of America Memoir 172, p. 405-424.

Jachens, R.C., Wentworth, C.M., and McLaughlin, R.J., 1998, Pre-San Andreas location of the Gualala block inferred from Magnetic and gravity anomalies, *in* Geology and tectonics of the Gualala block, northern California: SEPM Pacific Section Book 84, p. 27-64.

John, D.A., Wallace, A.R., Ponce, D.A., Fleck, R.B., and Conrad, J.E., 2000, New perspectives on the geology and origin of the northern Nevada Rift, *in* Cluer, J.K., Price, J.G., Struhsacker, E.M., Hardyman, R.F., and Morris, C.L., eds., *Geology and Ore Deposits 2000: The Great Basin and Beyond: Geological Society of Nevada Symposium Proceedings*, May 15-18, p. 127-154.

Kane, M.F., and Godson, R.H., 1985, Features of a pair of long-wavelength (>250 km) and short-wavelength (<250 km) Bouguer gravity maps of the United States, *in* Hinze, W.J., ed., *The utility of regional gravity and magnetic anomaly maps: 52nd annual international meeting of the Society of Exploration Monograph*, p. 46-61.

Karlstrom, K.E., Bowring, S.A., Chamberlain, K R; Dueker, K.G., Eshete, T., Erslev, E.A., Farmer, G.L., Heizler, M., Humphreys, E.D., Johnson, R.A., Keller, G.R., Kelley, S.A., Levander, A., Magnani, M.B., Matzel, J.P., McCoy, A.M., Miller, K.C., Morozova, E.A., Pazzaglia, F.J., Prodehl, C., Rumpel, H.M., Shaw, C.A., Sheehan, A.F., Shoshitaishvili, E., Smithson, S.B., Snelson, C.M., Stevens, L.M., Tyson, A.R., and Williams, M.L., 2002, Structure and evolution of the lithosphere beneath the Rocky Mountains; initial results from the CD-ROM experiment, Continental Dynamics of the Rocky Mountains Working Group, United States: *GSA Today*, v. 12, p. 4-10.

Kleinhampl, F.H., and Ziony, J.I., 1984, Mineral resources of northern Nye County, Nevada: Nevada Bureau of Mines and Geology Bulletin 99B, 243p.

Kleinhampl, F.J., Davis, W.E., Silberman, M.L., Chesterman, C.W., Chapman, R.H., and Gray, C.H., Jr., 1975, Aeromagnetic and limited gravity studies and generalized geology of the Bodie Hills region, Nevada and California: U. S. Geological Survey Bulletin 1384, 38p.

Kuntz, M.A., Champion, D.E., Lefebvre, R.H., and Covington, H.R., 1988, Geologic map of the Craters of the Moon, Kings Bowl, and Wapi lava fields, and the Great Rift volcanic rift zone, south-central Idaho: U. S. Geological Survey Miscellaneous Investigations Series I-1632.

Kuntz, M.A., Covington, H.R., and Schorr, L.J., 1992, An overview of basaltic volcanism of the eastern Snake River plain, Idaho: Geological Society of America Memoir 179, p. 227-267.

Lachenbruch, A.H., and Sass, J.H., 1978, Heat flow in the United States and the thermal regime of the crust: U. S. Geological Survey Professional Paper 1100, p. 200-201.



LaFehr, T.R., 1965, Gravity, isostasy, and crustal structure in the southern Cascade Range: Journal of Geophysical Research, v. 70, p. 5581-5597.

LaFehr, T.R., 1966, Gravity in the eastern Klamath Mountains, California: Geological Society of America Bulletin, v. 77, p. 1177-1190.

Langenheim, V.E., Glen, J.M.G., Jachens, R.C., Dixon, G.L., Katzer, T.C., and Morin, R.L., 2000, Geophysical constraints on the Virgin River Depression, Nevada, Utah, and Arizona: U.S. Geological Survey Open-File Report 00-0407.

Mabey, D.R., 1961, Regional magnetic and gravity anomalies in the Darwin area, California: U. S. Geological Survey Professional Paper 0424-C, p. C276-C279.

Mabey, D.R., 1976, Interpretation of a gravity profile across the western Snake River Plain, Idaho: Geology, v. 4, p. 53-55.

Mabey, D.R., 1982, Geophysics and tectonics of the Snake River plain, Idaho, *in* Bonnichsen, B., and Breckenridge, R.M., eds., Cenozoic geology of Idaho: Idaho Bureau of Mines and Geology Bulletin 26, p. 139-153.

Mabey, D.R., 1986, Gravity and magnetic anomaly maps of the Tenmile West Roadless Area, Boise and Elmore counties, Idaho: U. S. Geological Survey, Miscellaneous Field Studies Map MF-1500-D, 1 sheet.

Mabey, D.R., Zietz, I., Eaton, G.P., and Kleinkopf, M.D., 1978, Regional magnetic patterns in part of the Cordillera in the western United States, *in* Smith, R.B., and Eaton, G.P., eds., Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152, p. 93-106.

Mankinen, E.A., Hildenbrand, T.G., Dixon, G.L., McKee, E.H., Fridrich, C.J., and Laczniaik, R.J., 1999, Gravity and magnetic study of the Pahute Mesa and Oasis Valley region, Nye County, Nevada: U.S. Geological Survey Open-File Report 99-303, 35p.

McKee, E.H., 1968, Geology of the Magurder Mountain area, Nevada-California: U.S. Geological Survey Bulletin 1251-H, p. 1-40.

McKee, E.H., Duffield, W.A., and Stern, R.J., 1983, Late Miocene and early Pliocene basaltic rocks and their implications for crustal structures, northeastern California and south-central Oregon, *Geological Society of America Bulletin*, v. 94, p. 292-304.

McKee, E.H., Hildenbrand, T.G., Anderson, M.L., Rowley, P.D., and Sawyer, D.A., 1999, The Silent Canyon caldera complex - a three-dimensional model based on drill-hole stratigraphy and gravity inversion: U.S., Geological Survey Open-File Report 99-555, 38p.

McKee, E.H., Jachens, R.C., and Blakely, R.J., 1990, Major crustal differences between the northwestern part of the Great Basin and other parts of the province, *in* Cuffney, B., ed., *Geology and ore deposits of the Great Basin: Geological Society of Nevada and U.S. Geological Survey Program with abstracts*, p. 100.

McKee, E.H., and Nelson, C.A., 1967, Geologic map of the Soldier Pass Quadrangle, California and Nevada: U. S. Geological Survey Geologic Quadrangle Map Report GQ-0654.

McKee, E.H., Phelps, G.A., and Mankinen, E.A., 2001, The Silent Canyon caldera - a three dimensional model as part of a Pahute Mesa - Oasis Valley hydrogeologic model: U.S. Geological Survey Open-File Report 01-297.

Noble, D.C., and McKee, E.H., 1972, Description and K-Ar ages of volcanic units of the Caliente volcanic field, Lincoln County, Nevada, and Washington County, Utah: *Isochron/West*, v. 5, p. 17-24.

Oldow, J.S., 1984, Evolution of a late Mesozoic back-arc fold and thrust belt, northwestern Great Basin, U.S.A., *Tectonophysics*, 102, 245-274.

Oliver, H.W., 1977, Gravity and magnetic investigations of the Sierra Nevada Batholith, California: *Geological Society of America Bulletin*, v. 88, p. 445-461.

Oliver, H.W., 1980, Sierra Nevada, *in* Oliver, H.W., ed., *Interpretation of the gravity map of California and its continental margin: California Division of Mines and Geology Bulletin 205*, p. 30-32.

Oliver, H.W., and Griscom, A., 1980, Great Valley, *in* Oliver, H.W., ed., Interpretation of the gravity map of California and its continental margin: California Division of Mines and Geology, Bulletin 205, p. 27-29.

Oliver, H.W., and Mabey, D.R., 1963, Anomalous gravity field in east-central California, Geological Society of America Bulletin, v. 74, p. 1293-1298.

Oliver, H.W., Moore, J.G., and Sikora, R.F., 1992, Internal structure and vertical extent of the Sierra Nevada Batholith, California, from specific-gravity and gravity data, *in* Guangzhi, T., Keqin, Xu. And Yuzhou, Q., eds., Petrogenesis and mineralization of granitoids, Proceedings of 1987 Guangzhou International Symposium: Beijing, China, Science Press.

Oliver, H.W., Ponce, D.A., and Hunter, W. Clay, 1995, Major results of geophysical investigations at Yucca Mountain and vicinity, southern Nevada: U.S. Geological Survey Open-File Report 95-74, 184 p.

Page, B.M., 1965, Preliminary geologic map of a part of the Stillwater Range, Churchill County, Nevada: Nevada Bureau of Mines and Geology Map 28, scale 1: 125,000.

Pierce, K.L., Morgan, L.A., and Saltus, R.W., 2000, Yellowstone plume head: postulated tectonic relations to the Vancouver slab, continental boundaries and climate: U.S. Geological Survey Open-File Report 00-498, 39p.

Ponce, D.A., and Glen, J.M.G., 2002, Relationship of epithermal gold deposits to large-scale fractures in northern Nevada: Economic Geology, v. 97, p. 3-9.

Reford, M.S., 1980, Magnetic method, *in* Domenico, S.N., and Silverman, D., eds., Fifty years of progress; a review; 1930-1980: Geophysics, v. 45, p. 1640-1658.

Roberts, C.W., and Jachens, R.C., 1993, Isostatic residual gravity map of the San Francisco Bay area, California: U.S. Geological Survey Geophysical Investigations Map GP-1006.

Rowan, L.C., and Wetlaufer, P.H., 1973, Structural geologic analysis of Nevada using ERTS-1 images: a preliminary report: NASA Special Publication 327, p. 413-423.

Rytuba, J.J., and McKee, E.H., 1984, Perakaline ash-flow tuffs and calderas of the McDermitt volcanic field, southeast Oregon and north-central Nevada: *Journal of Geophysical Research*, v. 89, p. 9616-8628.

Saltus, R.W., Thompson, G.A., 1995, Why is it downhill from Tonopah to Las Vegas? A case for mantle plume support of the high northern Basin and Range: *Tectonics*, v. 14, p. 1235-1244.

Simpson, R.W., and Jachens, R.C., 1989, Gravity methods in regional studies, *in* Pakiser, L.C., Mooney, W.D., eds., *Geophysical framework of the continental United States*, Geological Society of America Memoir 172, p. 35-44.

Simpson, R.W., Jachens, R.C., Blakely, R.J., and Saltus, R.W., 1986, A new isostatic residual gravity map of the conterminous United States with a discussion on the significance of isostatic residual anomalies: *Journal of Geophysical Research*, v. 91, p. 8348-8372.

Sims, P.K., Bankey, V., and Finn, C.A., 2001, Preliminary Precambrian basement map of Colorado -- A geologic interpretation of the aeromagnetic anomaly map: U.S. Geological Survey Open-File Report 01-364.

Smith, R.B., 1978, Seismicity, crustal structure, and intraplate tectonics of the interior of the western Cordillera, *in* Smith, R.B., and Eaton, G.P., eds., *Cenozoic tectonics and regional geophysics of the Western Cordillera*: Geological Society of America Memoir 152, p.111-144.

Sparlin, M.A., Braile, L.W., and Smith, R.B., 1982, Crustal structure of the eastern Snake River plain determined from ray trace modeling of seismic refraction data: *Journal of Geophysical Research*, v. 87, p. 2619-2633.

Speed, R.C., 1976, Geologic map of the Humboldt lopolith: Geological Society of America Map and Chart Series MC-14, scale 1:81,050.

Stewart, J.H., 1988, Tectonics of the Walker Lane belt, western Great Basin: Mesozoic and Cenozoic deformation in a zone of shear, *in* Ernst, W.G., ed., *Metamorphism and crustal evolution of the western United States*, Rubey volume VII: Englewood Cliffs, New Jersey, Prentice Hall, p. 683-713.

Stewart, J.H., and Carlson, J.E., 1978, Geologic map of Nevada: Nevada Bureau of Mines and Geology Map, scale 1:500,000.

Stewart, J.H., and Crowell, J.C., 1992, Strike-slip tectonics in the Cordilleran region, Western United States, *in* Burchfiel, B.C., Lipman, P.W., and Zoback, M.L., The geology of North America, the Cordilleran Orogen; conterminous U.S.: Boulder, Colorado: Geological Society of America, v. G-3, p. 609-628.

Stewart, J.H., Moore, W.J., and Zietz, I., 1977, East-west patterns of Cenozoic igneous rocks, aeromagnetic anomalies, and mineral deposits, Nevada and Utah: Geological Society of America Bulletin, v. 88, p. 67-77.

Sumner, J.S., 1985, Crustal geology of Arizona as interpreted from magnetic, gravity, and geologic data, *in* Hinze, W.J., ed., The utility of regional gravity and magnetic anomaly maps: 52nd annual international meeting of the Society of Exploration Monograph, p. 164-180.

Suppe, J., 1979, Structural interpretation of the southern part of the northern Coast Ranges and Sacramento Valley, California: Geological Society of America Bulletin, v. 90, p. I327-I 330.

Thompson and McCarthy, 1990, A gravity constraint on the origin of highly extended terranes: Tectonophysics, v.174, p.197-206.

U. S. Geological Survey, 1972, Aeromagnetic map of the Vya and part of the McDermitt 1 degrees by 2 degrees quadrangles, Nevada, U.S. Geological Survey Open-File Report 72-0393.

U. S. Geological Survey, 1982a, Spectral radiometric and total-field magnetic survey of the McDermitt calderas, NV-OR: U. S. Geological Survey Open-File Report 82-0323-A, 48p.

U. S. Geological Survey, 1982b, Stacked magnetic and radiometric profiles of flight lines in the McDermitt calderas area, Nevada-Oregon: U.S. Geological Survey Open-File Report 82-0323-B, 96p.

Vallier, T.L., and Brooks, H.C., eds., 1986, Geology of the Blue Mountains region of Oregon, Idaho and Washington: Geologic implications of Paleozoic and Mesozoic paleontology and biostratigraphy, Blue Mountains Province, Oregon and Idaho: U.S. Geological Survey Professional Paper 1435, 93p.

Vallier, T.L., and Brooks, H.C., eds., 1994, Geology of the Blue Mountains region of Oregon, Idaho and Washington: Stratigraphy, physiography and mineral resources of the Blue Mountains region: U.S. Geological Survey Professional Paper 1439, 198p.

Vallier, T.L., and Brooks, H.C., eds., 1995, Geology of the Blue Mountains region of Oregon, Idaho and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region: U.S. Geological Survey Professional Paper 1438, 540p.

Vogel, T.A., Cambray, F.W., and Constenius, K.N., 2001, Origin and emplacement of igneous rocks in the central Wasatch Mountains, Utah: *Rock Mountain Geology*, v. 36, p. 119-162.

Wells, R.E., Weaver, C.S., and Blakely, R.J., 1998, Fore-arc migration in Cascadia and its neotectonic significance: *Geology*, v. 26, p. 759-762.

Woollard, G.P., 1972, Regional variations in gravity, *in* Robertson, E.C., ed., *The Nature of the Solid Earth*: New York, McGraw-Hill, p. 463-505.

Wyld, S.J., and Wright, J.E., 2001, New evidence for Cretaceous strike-slip faulting in the United States Cordillera and implications for terrane-displacement, deformation patterns, and plutonism: *American Journal of Science*, v. 301, p. 150-181.

Zietz, I., Bateman, P.C., Case, J.E., Crittenden, M.D., Jr., Griscom, A., King, E.R., Roberts, R.J., and Lorentzen, G.R., 1969, Aeromagnetic investigation of crustal structure for a strip across the western United States: *Geological Society of America Bulletin*, v. 80, p. 1703-1714.

Zoback, M.L., and Thompson, G.A., 1978, Basin and range rifting in northern Nevada; clues from a mid-Miocene rift and its subsequent offsets: *Geology*, v. 6, p. 11-116.

Zoback, M.L., and Zoback, M.D., 1980, Magnitude of deviatoric stresses in the Earth's crust and uppermost mantle: *Journal of Geophysical Research*, v. 85, p. 6113-6156.

Zoback, M.L., McKee, E.H., Blakely, R.J., and Thompson, G.A., 1994, The northern Nevada rift; regional tectono-magmatic relations and middle Miocene stress direction: *Geological Society of America Bulletin*, v. 106, p. 371-382.

## **TABLES**

Table 1. Geophysical features of the Great Basin and parts of surrounding provinces. Terranes appear in the table sorted in the order of: state, gravity number, and magnetic number. Refer to gravity and magnetic terrane maps (figs. 2 and 3, respectively). The table gives (from left to right): Scale of anomaly (R=Regional, I=Intermediate, L=Local); Terrane ID (containing from left to right: 2 digits for the state, 1 'm' or 'g' for magnetic or gravity anomaly, respectively; two digits for terrane number); Terrane definition (describes the geophysical characteristics that define the feature), Anomaly type (H=high, L=low, or B=both high and low); Geologic Province (describes the geology associated with the geophysical terrane); Inferred source of anomaly; Generalized source rock (see Table 3 for description of terms); Tectonic setting (see table 4 for description of terms); Province (gives the associated physiographic province, see table 2); References. Abbreviations include: BR, Basin and Range; CP, Colorado Plateau; CRP, Columbia River Plateaus; Cz, Cenozoic; GB, Great Basin; Mz, Mesozoic; NNR, Northern Nevada Rift; pC, Precambrian; Pz, Paleozoic; SRP, Snake River Plain. For further details, see text.

Table 2. Physiographic provinces and their geologic and geophysical character. Physiographic province boundaries and descriptions are modified from Fenneman and Johnson (1946).

## **FIGURES**

Figure 1. A) Index map of the Great Basin showing digital shaded relief and outlines of the Great Basin and surrounding physiographic provinces. Physiographic province boundaries are modified from Fenneman and Johnson (1946); B) The Basin and Range extensional province that includes the Great Basin.


Figure 2. Basement gravity terrane map of the study area. Blue lines are edges of gravity sources determined by the maximum horizontal gravity method; each feature is uniquely labeled and described in Table 1. Yellow lines are linear gravity features, labeled uniquely and described in Table 1; red lines are unlabeled, undescribed, linear gravity features. White line

shows boundary within which basement calculation was performed. Isostatic gravity is shown outside this boundary.

Figure 3. Magnetic potential terrane map of the study area. Blue lines are edges of magnetic sources determined by the maximum horizontal gravity method; each feature is uniquely labeled and described in Table 1. Yellow lines are linear magnetic features, labeled uniquely and described in Table 1; red lines are unlabeled, undescribed, linear magnetic features.

Figure 4. Regional isostatic gravity and pseudogravity field maps of the Great Basin and surrounding area. The white outline represents the Great Basin boundary in each map. The black outlines represent the regional gravity terrane and regional pseudogravity terrane described in Table 1 as G1 and M1, respectively.








Figure 2. Basement gravity terrane map of the Great Basin. Blue lines circumscribe labeled polygonal features described in Table 1; yellow lines indicate labeled linear features described in Table 1; red lines indicate undescribed linear trends.






Figure 3. Magnetic potential (pseudogravity) terrane map of the Great Basin. Blue lines circumscribe labeled polygonal features described in Table 1; yellow lines show labeled linear features described in Table 1; red lines show undescribed linear trends.




Figure 4. Regional isostatic gravity and pseudogravity field maps of the Great Basin.